



Universitas Gadjah Mada
Institute for Halal Industry & System

ISBN : 978-602-17639-1-9

Proceeding

The 3rd International Symposium toward Halal Global 2019

10 Desember 2019



Proceeding

The 3rd International Symposium Toward Halal Global

The contributions of Scientist toward Halal Global

Yogyakarta, December 10, 2019
University Club Hotel, Universitas Gadjah Mada

ISBN:
978-602-17639-1-9

Publisher
Laboratorium Penelitian dan Pengujian Terpadu (LPPT)
UNIVERSITAS GADJAH MADA

Proceeding
the 3rd International Symposium Toward Halal Global

The contributions of Scientist toward Halal Global

Organizing Committee

Chairman : Prof. Ir. Yuny Erwanto, S.Pt., MP., Ph.D.

Secretary : Dr. Lily Arsanti Lestari, STP., MP.

Treasurer : M. Fahmi Dwi Ari Yuwono, SE.

Secretariat : Tri Cahya Wibawa, S.Kom.

Fitriana Maharjanti, S.Si.

Scientific Committee:

1. Ir. Yuny Erwanto, S.Pt., MP., Ph.D.
2. Dr. Lily Arsanti Lestari, STP., MP.
3. Prof. Dr. Abdul Rohman, M.Si., PhD., Apt.
4. Tri Joko Raharjo, Ph.D.
5. Dr. Kuwat Triyana

Supporting Organizing Committee (students)

1. Deny Hermawan
2. Fredo Wahyu Ramanda
3. Muhammad Faris Al Akram
4. Muhammad Ridwan
5. Nisa Muflichati
6. Siti Aisyah
7. Dewi Sarah
8. Siska Mardiana
9. Nadya Ramadhani Susilo

Table of Content

Organizing Committee		ii	
Preface		iv	
Table of Content		v	
Extended Abstract of Invited Speaker			
1. Halal certification in South Korea and collaboration opportunities (Dr. James Noh)		1	
2. National Halal Market Potency & Strategy to Penetrate Global Market (Akhmad Akbar Susanto, SE., M.Phil., PhD)		4	
3. The role of standardized analytical methods in supporting halal certification in Indonesia (Prof. Dr. Abdul Rohman, M.Si., Apt.)		5	
4. Real-Time PCR Technology for Halal Authentication (Assoc. Prof. Dr. Kartini Abu Bakar)		7	
Abstract of Poster Presenter			
P-IE-1	Does Sharia Governance Exist? Evidence on Islamic Banking in Indonesia	Ely Windarti Hastuti	12
P-IE-2	Virtual Reality for Islamic Financial Literacy: A Shift in the New Education Strategy for the Digital Generation	Ratna Candra Sari	12
P-IE-3	Designing Virtual Assistant for Islamic Financial Literacy Using Augmented Reality Technology	Sariyatul Ilyana	13
P-HL-1	Creating Bangunjiwo as a Halal Food Tourism Village	Ambar Rukmini	13
P-HL-2	Halal Tourism Index (HTI) and Model of Halal Tourism Management in Bukittinggi City, West Sumatera	Muhammad Ghafur Wibowo	14
P-HL-3	The role of Accounting Information Systems in Halal Food Supply Chain.	Donny Setiawan	14
P-HL-4	Should Halal Certification Be a Bad Product Brand Connotation? An Islamic And Positive Juridical Review	Ridan Muhtadi	15
P-HL-5	Muslim Student's Awareness of Halal Food Products Small Micro Entrepreneurs	Hamii Datur RosyDAH	15
P-HL-6	Social Awareness in Checking Halal Labels Before Ordering Food Online	Durrotun Nafisah	16
P-HL-7	New Start-Up Development Model "Halalin Aja" Using Business Model Canvas	Anggun Fambudi Ningrum Lestari	16
P-HPDD-1	The Synthesis and Application of Edible Film for Eco-Friendly Material of Halal Packaging	Neneng Windayani	17
P-HPDD-2	Application of FTIR Spectroscopy for Initial Analysis of Leather Products Adulteration of Pig Skin	Ragil Yuliatmo	18

P-HPDD-3	Utilization of <i>Rhizopus Oryzae</i> on The Development of Halal Dairy Product Based on Biotechnology	Solikah Ana Estikomah	18
P-HPDD-4	Porcine DNA Contamination Test on Meat Milling Sample Using the Real Time PCR Method	Yuanita Rachmawati	19
P-FSN-1	In Vitro Test: Screening the Effectiveness of Reducing Glucose Levels of Parijoto Extract the Chitosan Encapsulated	Fania Putri Luhurningtyas	19
P-FSN-2	Screening of Antioxidant and Reducing Glucose Activity of Arabica Green Coffee Bean (<i>Coffea Arabica</i> L.) Purified Extract by UV-Vis Spectrophotometry	Rissa Laila Vifta	20
P-FSN-3	Intake a Mixture of Halal Lamb Meat and Fat that was Roasted can Increase Blood Glucose Levels	Fathimah	20
P-FSN-4	Antioxidant Activities of <i>Strobilanthes crispus</i> Leaf Extracts in Streptozotocin-induced Diabetic Rats	Veriani Aprilia	21
 FullPaper			
1	Does Sharia Governance Exist? Evidence on Islamic Banking in Indonesia (Ely Windarti Hastuti, Fajar Amtsal, Siti Lisnawati)		23
2	Should Halal Certification Be a Bad Product Brand Connotation? an Islamic and Positive Juridical Review (Ridan Muhtadi, Mohsi, Kamali)		29
3	Muslim Student's Awareness of Halal Food Products Small Micro Entrepreneurs (Hamii Datur Rosyidah, Durrotun Nafisah, Hafidhotun Nabawiyah)		35
4	Social Awareness in Checking Halal Labels Before Ordering Food Online (Durrotun Nafisah, Hamii Datur Rasydah, Hafidhotun Nabawiyah)		39
5	Initial Analysis of Pigskin Adulteration on Leather Products Using FTIR Spectroscopy (Ragil Yuliatmo, Wisnu Pambudi, Thoyib Rahman Hakim, R.L.M.S. Ari Wibowo, Dwi Wulandari, and Yuny Erwanto)		43
List of Participant			51

Full Paper 5

INITIAL ANALYSIS OF PIGSKIN ADULTERATION ON LEATHER PRODUCTS USING FTIR SPECTROSCOPY

Ragil Yuliatmo^{1*}, Wisnu Pambudi², Thoyib Rahman Hakim¹, R.L.M.S. Ari Wibowo¹,
Dwi Wulandari¹, and Yuny Erwanto³

¹) Department of Leather Processing Technology, Politeknik ATK Yogyakarta,
Yogyakarta, Indonesia email: info@atk.ac.id

²) Department of Rubber and Plastic Processing Technology, Politeknik ATK Yogyakarta,
Yogyakarta, Indonesia email: info@atk.ac.id

³) Department of Animal Product Technology, Faculty of Animal Science, Universitas
Gadjah Mada, Yogyakarta, Indonesia, email: fapet@ugm.ac.id

* Corresponding Author: E-mail: ragilyuliatmo@kemenperin.go.id

ABSTRACT

Leather products are part of daily fashion in Indonesia, such as bags, shoes, jackets, gloves, etc. Adulteration of raw materials for leather products can occur if there are no labels on these products. The existence of certain religion / belief that prohibit adherents from using certain materials, such as pork skin. Detection of adulteration has been carried out in various ways such as by PCR, GC-MS, HPLC, and FTIR methods. The FTIR method is known as an easy and inexpensive method to use. The objective of this study was to evaluate the capability of FTIR spectroscopy for lipid identification and initial analysis to detection of pig skin on leather products. Lipid extracts obtained from the various skin were scanned using FTIR spectrophotometer at 4000–500 cm^{-1} . It resulted spectral differences in several wavenumber (2951-3258 cm^{-1} and 1046-1428 cm^{-1}). At wavenumber 3020-2980 cm^{-1} there is a peak only in lard. The same result is also found in lipid spectra from leather products extraction. The FTIR spectroscopy is able to differentiate pigskin from goat and sheep skins through specific peaks in infrared spectra. This can be used as an initial analysis on determining the existence of skin adulteration in leather products. This study is prospective to be continued by chemometrics as a quantitative analysis

Keywords: Adulteration, FTIR Spectroscopy, Initial Analysis, Leather Products, Pigskin

1. INTRODUCTION

Determination of authentication and detection of adulteration of material products is one of the main issues in the industrial field (Marikkar et al., 1995; Al Jowder et al., 1997). One of the risky products to be falsified is leather products. Detection of product adulteration is important for consumer protection and also for certain religious reasons. In some countries, producers and sellers of leather products choose to use pigskin as a substitute for other skins, because prices are cheap and easy to obtain (Aida et al., 2005). The importance of labeling is influenced by the existence of certain religions / beliefs that prohibit adherents from using certain ingredients, such as the prohibition of the use of the element of pigs for Muslim and Jewish communities (Nakyisinge et al., 2012). Recently, the inclusion of material of animal origin is often not displayed clearly so scientific proof is needed to find out.

Various techniques or methods have been used for pig elemental analysis, such as Gas Chromatography-Mass Spectroscopy (Nizar et al., 2013), Liquid Chromatography-Mass Spectroscopy (LC-MS) (Czerwenka et al., 2010), Gas Chromatography Tandem Mass Spectrometry (GC-MS) (Oliveira et al., 2009), Differential Scanning Calorimetric (DSC) (Marina et al., 2009; Nurrulhidayah et al., 2015), High Performance Liquid Chromatography (HPLC) (Saeed et al., 1989; Marikkar et al., 2005), Electronic Nose

(Nurjuliana et al., 2011), and DNA-based methods using the polymerase chain reaction (PCR) method (Che Man et al., 2011; Erwanto et al., 2014; Maryam et al., 2015). Some methods that have been done have weaknesses because it requires a long time and expensive costs in detecting adulteration in food. Therefore, routine methods that are fast, accurate, inexpensive, and easy to use are needed. One ideal method to be used in laboratories is Fourier Transform Infrared (FTIR) spectroscopy. Application of FTIR for initial analysis on leather products is appropriate to be studied.

2. METHODS

Lipid extraction

Lipid extraction using Soxhlet method was performed according to AOAC (1995). Raw leather and leather products in the form of leather are obtained from the market and leather distributors. A-50.0 g of samples was wrapped with filter paper and placed into the Soxhlet apparatus. A-250 mL of n-hexane was used as extracting solvent. The extraction was performed for 8 h at 70°C (± 50 cycles). The lipid extract was added with anhydrous sodium sulfate, mixed, filtered by filter paper, and then evaporated until the solvent was completely removed. The resulting lipid fraction is then used for FTIR spectral measurements.

FTIR spectral measurements

The lipids obtained by lipid extraction were placed in attenuated total reflectance (ATR) crystal at ambient temperature (25°C). The spectrum was acquired in the wavenumbers region of 450–4000 cm^{-1} using FTIR spectrophotometer (Perkin-Elmer, Singapore).

Data analysis

The spectrum from FTIR spectral measurement results were analyzed descriptively by comparing of several skin species from raw skin and leather

3. RESULTS AND DISCUSSION

Lipid extraction

Raw material skin and leather have extracted by sokhlet method, and the solvent used by n-hexane. N-hexane solvent is very suitable for use in the extraction of fat on the skin. This is because n-hexane is non-polar, just like fat which is also non-polar and n-hexane is easily evaporated because it has a low boiling point of 69°C (Erwanto et al, 2016). The extraction of fat from all types of species produced yellowish and thick oil, but specifically in sheep skin fat produces oil that is easily frozen at room temperature, while others do not.

FTIR Analysis of fresh skin lipid

The FTIR spectra of lipid obtained from four species skin have similar profiles. Figure 1 showed lipid spectra of pig, goat, sheep, and cattle skin lipid. Infrared spectra is read at 4000-550 cm^{-1} , which is the middle region. Many molecules have a strong absorbance in the middle infrared region. Many types of samples including solids, liquids, gases, semisolids, powders, polymers, organic, inorganic, biological substances, pure substances, and mixtures can be measured in the middle region (Smith, 2011).




Figure 1. FTIR spectra of the lipid extracted from pig, goat, sheep, and cattle skin.

The results showed at a glance that there is no difference in the peak point in each fat, but it observed in more detail, there are some different peak point in the absorption of functional groups. There are at least 2 different spectra between each of the sources $2951\text{-}3258\text{ cm}^{-1}$ (a) and $1046\text{-}1428\text{ cm}^{-1}$ (b).

FTIR Analysis of lipid from leather products

Figure 2 showed that the spectra of three leather products have characters that are almost the same as the spectra of raw skin.

Goat fat

Sheep fat




Figure 2. FTIR spectra of the lipid extracted from pig, goat, and sheep leather

Based on the results in Figure 2, wavenumber of the spectra is enlarged to be in the range of 3000-2800 cm^{-1} . The magnification of the area is shown in Figure 3. In that range, there is a peak in the spectra of pig skin leather lipid, while other types of leather do not have peaks in the wavenumber, this is also found in the raw skin spectra.

FTIR Analysis of lipid from commercial fatliquoring agent

In softy leather, as in leather products in this study, the addition of a fatliquor agent will improve the leather quality. Fatliquoring used fatliquor agent which is generally derived from the fat element, so that it can affect the spectra on crust skin (Covington, 2009). Fatliquoring is one of the important stages in the final process of tanning the skin so that it gets the desired leather character, such as the soft character on the glove or the medium hard on lining or upper shoes (Zarlok et al., 2014). This is thought to affect the fat spectra on the leather.




Figure 3. The enlarged FTIR spectra at wavenumber regions 3000-2800 cm^{-1}

Three types of fatliquoring agents or commercial fatliquors have been analyzed by FTIR in Figure 4. In the range of wavenumber 3700-3000 cm^{-1} there is a peak which showed the group H = O. This can be interpreted that the fatliquor still contains water (H_2O), while the main source of fatliquor is fat, so it is suspected that the three fatliquors contain surfactants that can bind fat and also water at the same time.




Figure 4. FTIR spectra of the lipid extracted from fatliquoring agent (Catalyc GS, Dermal ESN, dan Pelan GLH)

In wavenumber 300-2800 cm^{-1} there is a peak which has almost the same character in the leather fat spectra. Figure 5 showed the magnification of the peak of the fat spectra. Based on these results it was found that the wavenumber had similarities with the spectra of goat and sheep crust skin although the absorbance of fatliquor was relatively higher, but did not have the same characteristics as pig leather.




Figure 5. The enlarged FTIR spectra of commercial fatliquoring agents at wavenumber regions 3000-2800 cm^{-1} .

4. CONCLUSION

Lipid from skin extracts tested by FTIR have produced spectral differences in several wavenumber ($2951\text{-}3258\text{ cm}^{-1}$ and $1046\text{-}1428\text{ cm}^{-1}$). At wavenumber $3020\text{-}2980\text{ cm}^{-1}$ there is a peak only in lard. The same thing is also found in lipid spectra from leather products extraction. The addition of fatliquor did not affect the specificity of pig skin rind spectra. The FTIR spectroscopy is able to differentiate pigskin from goat and sheep skins through specific peaks in infrared spectra. This can be used as an initial analysis on determining the existence of skin adulteration in leather products. This study is prospective to be continued by chemometrics as a quantitative analysis

5. ACKNOWLEDGEMENT

The authors would like to thank the Research and Community Service Unit in Politeknik ATK Yogyakarta for financial support via Research Project 2019

REFERENCES

- Aida, A.A., Y.B.C. Man, C.M.V.L. Wong, A.R. Raha and R. Son, 2005. Analysis of raw meats and fats of pigs using polymerase chain reaction for Halal authentication. *Meat Sci.*, 69: 47-52.
- Al-Jowder, O. Kemsley, E.K. Wilson, R.H. 1997. Mid-infrared spectroscopy and authenticity problems in selected meats: a feasibility study. *Food Chem.*, 59, 195-201.
- AOAC (Association of Official Analytical Chemists). 1995. Official Methods of Analysis. AOAC International, Gaithersburg, MD. 16th Edition
- Covington, D. 2009. Tanning Chemistry: The Science of Leather. The Royal Society of Chemistry. Cambridge, UK.
- Czerwenka, C., L. Muller dan W. Lindner. 2010. Detection of the adulteration of water buffalo milk and mozzarella with cow's milk by liquid chromatography-mass spectrometry analysis of β -lactoglobulin variants. *Food Chem.*, 122: 901-908
- Che Man, Y.B. Rohman, A. Mansor, T.S.T. 2011. Differentiation of lard from other edible oils by means of Fourier transform infrared spectroscopy and chemometrics. *J Amer Oil Chemists' Soc.*, 88,187–192.
- Erwanto, Y. Abidin, M.Z. Muslim, E.Y.P. Sugiyono, Rohman. 2014. A. Identification of pork contamination in meatballs of Indonesia local market using polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) analysis. *Asian Aus. J. Anim Sci*, 27: 1487-1492.
- Erwanto, Y. A.T. Muttaqien, Sugiyono, Sismindari, A. Rohmanet. 2016. Use of Fourier Transform Infrared (FTIR) Spectroscopy and Chemometrics for Analysis of Lard Adulteration in "Rambak" Crackers. *International J Food Properties*. Taylor & Francis, 19(12), pp. 2718–2725. doi: 10.1080/10942912.2016.1143839
- Marikkar, J.M.N.; Ghazali, H.M.; Che Man, Y.B.; Peiris, T.S.G.; Lai, O.M. 2005. Distinguishing lard from other animal fats in admixtures of some vegetable oils using liquid chromatographic data coupled with multivariate data analysis. *Food Chem.*, 91, 5–14.
- Marina, A.M.; Che Man, Y.B.; Nazimah, S.A.H.; and Ismail, A. 2009. Monitoring the adulteration of virgin coconut oil by selected vegetable oils using differential scanning calorimetry. *J. Food Lipids*, 16:50–61
- Maryam, S.; Sismindari; Raharjo, T.J.; Sudjadi; Rohman, A. 2005. Analysis of porcine contamination in dendeng using mitochondrial D-loop 686 and cyt B gene primers by real time polymerase chain reaction.
- Nakyinsige, K., Man, Y. B. C., & Sazili, A. Q. (2012). Halal authenticity issues in meat and meat products. *Meat Sci.*, 91(3), 207–214. doi:10.1016/j.meatsci.2012.02.015.
- Nizar, N.N.A.; Marikkar, J.M.N.; Hashim, D.M. 2013. Differentiation of lard, chicken fat, beef fat and mutton fat by GCMS and EA-IRMS techniques. *J Oleo Sci.*, 62, 459-464.